



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN ALL STATES EXCEPT NEW YORK AND TEXAS

HYDROTHOL® GRANULAR AQUATIC ALGICIDE AND HERBICIDE EPA Reg. No. 70506-174

FOR CONTROL OF AQUATIC PLANTS LISTED ON THE TABLE BELOW

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

RATE OF APPLICATION

| Aquatic Plants | Rate (ppm) | Hydrothol Granular |
|--|--------------|--------------------|
| | | Lbs/Ac.Ft. |
| Coontail, <i>Ceratophyllum</i> spp. | 1.5-2.5 ppm | 80-135 lbs |
| Horned Pondweed, <i>Zannichellia palustris</i> | 1.5-2.5 ppm | 80-135 lbs |
| Milfoil (<i>Myriophyllum</i> spp) | 1.5-2.5 ppm | 80-135 lbs |
| Sago Pondweed, <i>Stuckenia pectinata</i> | 1.0-2.5 ppm | 55-135 lbs |
| Naiad (<i>Najas</i> spp) | 1.5-2.5 ppm | 80-135 lbs |
| Pondweed (<i>Potamogeton</i> spp) Potamogeton spp. | 0.75-2.5 ppm | 40-135 lbs |
| Including: | | |
| American, <i>P. nodosus</i> | 1.5-2.5 ppm | 80-135 lbs |
| Largeleaf (Bass Weed), <i>P. ampifolius</i> | 1.5-2.5 ppm | 80-135 lbs |
| Curlyleaf, <i>P. crispus</i> | 0.75-1.5 ppm | 40-80 lbs |
| Flatstem, <i>P. zosteriformis</i> | 1.5-2.5 ppm | 80-135 lbs |
| Floating-leaf, <i>P. natans</i> | 1.0-2.5 ppm | 55-135 lbs |
| Illinois, <i>P. illinoensis</i> | 1.25-2.5 ppm | 65-135 lbs |
| Narrowleaf, <i>P. pusillus</i> | 1.0-2.5 ppm | 55-135 lbs |
| Threadleaf, <i>P. filiformis</i> | 1.5-2.5 ppm | 80-135 lbs |
| Variable Leaf, <i>P. diversifolius</i> | 1.0-2.5 ppm | 55-135 lbs |
| Water Stargrass* | 1.5-2.5 ppm | 80-135 lbs |

*Not for use in California